



King's Research Portal

DOI:

[10.1017/S0033291714001354](https://doi.org/10.1017/S0033291714001354)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Munafò, M. R., & Kempton, M. J. (2015). Has analytical flexibility increased in imaging studies of bipolar disorder and major depression? *Psychological Medicine*, 45(3), 449-451. <https://doi.org/10.1017/S0033291714001354>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Post print version – final PDF can be found at <http://dx.doi.org/10.1017/S0033291714001354>

Running Head: Analytical Flexibility in Imaging Studies

Has Analytical Flexibility Increased in Imaging Studies of Bipolar Disorder and
Major Depression?

Marcus R. Munafò ^{1,2}, Matthew J. Kempton ³

1. UK Centre for Tobacco and Alcohol Studies and School of Experimental Psychology, University of Bristol, Bristol, United Kingdom.
2. MRC Integrative Epidemiology Unit (IEU) at the University of Bristol, Bristol, United Kingdom.
3. Department of Neuroimaging, Institute of Psychiatry, King's College London, London, United Kingdom

Corresponding Author: Marcus R. Munafò PhD, School of Experimental Psychology, University of Bristol, 12a Priory Road, Bristol BS8 1TU, United Kingdom. T: +44.117.9546841; F: +44.117.9288588; E: marcus.munaf@bristol.ac.uk

Abstract

There has been extensive discussion of problems of reproducibility of research. Analytical flexibility may contribute to this, by increasing the likelihood that a reported finding represents a chance result. We explored whether analytical flexibility has increased over time, using human imaging studies of bipolar disorder and major depression. Our results indicate that the number of measures collected per study has increased over time for studies of bipolar disorder, but not for studies of major depression.

Keywords: Bipolar Disorder; Major Depression; Imaging; Analytical Flexibility.

Has Analytical Flexibility Increased in Imaging Studies of Bipolar Disorder and
Major Depression?

There has been extensive discussion of problems of reproducibility of research across a range of scientific disciplines (Ioannidis, 2005). A number of factors have been identified which may contribute to this, such as data fabrication (Simonsohn, 2013), publication bias (Smulders, 2013), peer review methods (Park *et al.*, 2013) and low statistical power (Button *et al.*, 2013). For the most part these are not new concerns; however, one factor which may have changed over recent years is the scope for flexible data analysis, given the increasing automation of statistical analyses, and the ease with which multiple outcomes can be tested in the same data set.

The impact of flexible analytical procedures has recently been described by Simmons and colleagues (Simmons *et al.*, 2011), who conclude that it is “unacceptably easy to accumulate (and report) statistically significant evidence for a false hypothesis”. This problem is not confined to behavioural experiments in psychology – Carp recently reviewed 241 functional MRI studies and showed that there were almost as many unique analytical pipelines reported as there were individual studies, with many studies not reporting critical methodological details (Carp, 2012). However, it is not clear whether analytical flexibility has increased over time.

We therefore investigated whether analytical flexibility in structural imaging studies of bipolar disorder and major depression has increased over time, using the number of measures collected as a proxy index of analytical flexibility. With more measures available, there is greater scope for conducting multiple statistical tests, and

selecting those which provide the clearest results for reporting or highlighting. We also investigated whether the number of participants tested has increased over time.

Data were taken from the Bipolar Disorder Neuroimaging database (bipolardatabase.org) (Kempton *et al.*, 2008) and the Major Depression Neuroimaging Database (depressiondatabase.org) (Kempton *et al.*, 2011). These online databases include peer-reviewed computerised tomography and structural MRI studies which compare patients with bipolar disorder or patients with major depression, diagnosed using standard diagnostic criteria, to a healthy control group. From studies within these two databases, the total number of participants (patients + controls) and total number of different brain measures recorded per study were extracted for the present analyses. Brain measures were defined as the measurement of a brain region (e.g., left hippocampus volume) or the measurement of a cerebral abnormality (e.g., the presence of periventricular hyperintensities).

We used linear regression to explore the relationship between year of publication, number of measures and number of participants. For studies of bipolar disorder ($k = 141$), year of publication was not associated with number of participants ($B = -0.01$, 95% CI -0.02 to $+0.01$, $R^2 = 0.01$, $P = 0.23$) but was positively associated with number of measures ($B = +0.04$, 95% CI $+0.02$ to $+0.06$, $R^2 = 0.07$, $P = 0.001$). However, for studies of major depression ($k = 225$), year of publication was positively associated with number of participants ($B = +0.02$, 95% CI $+0.01$ to $+0.03$, $R^2 = 0.04$, $P = 0.001$) but not with number of measures ($B = +0.01$, 95% CI -0.01 to $+0.03$, $R^2 = 0.01$, $P = 0.21$). A Z-test indicated that these estimates differed, with strong evidence for number of participants ($P = 0.004$) and weaker evidence for number of measures ($P = 0.080$). These results are shown in Figure 1.

Insert Figure 1 about here.

Our results partly support the possibility that analytical flexibility has increased over time. Among structural imaging studies of bipolar disorder, the number of measures taken per study (assumed here to be a proxy index of analytical flexibility) has increased, while the average sample size has not. However, among structural imaging studies of major depression we observed the opposite pattern, with no increase in the number of measures taken per study but an increase in average sample size. The reasons for this discrepancy are not clear. We restricted our analysis to structural MRI region of interest studies because relevant analysis techniques are well-established, and therefore consistent across studies. While strength of the MRI scanner and slice thickness may influence results, we previously found no evidence that these factors influenced measures of six key brain regions (Kempton *et al.*, 2011). It is possible that our results represent chance findings, but the statistical evidence is sufficiently strong that this explanation is unlikely. The results also do not appear to be driven by a small number of outliers.

One possibility is that there are in fact fewer true effects in bipolar disorder compared to major depression (or the effects are considerably smaller). If it is harder to detect effects this may lead to increased pressure to collect multiple measures to increase the likelihood of finding something. The addition of future study databases recording analytical flexibility may clarify the apparent discrepancy between the major depression and bipolar disorder literatures. More generally, there is growing interest in methods to interrogate published literature. Our approach, which uses number of measures as a metric of analytical flexibility, may be useful as a scalable tool for analyzing all available studies across a published literature.

Acknowledgements

MRM is a member of the United Kingdom Centre for Tobacco and Alcohol Studies, a UKCRC Public Health Research: Centre of Excellence. Funding from British Heart Foundation, Cancer Research UK, Economic and Social Research Council, Medical Research Council, and the National Institute for Health Research, under the auspices of the UK Clinical Research Collaboration, is gratefully acknowledged. MJK is supported by and MRC Career Development Fellowship (grant number MR/J008915).

Author Contributions

MRM conceived of the study. MJK collected the data. Both authors contributed to analysis of the data and the writing of the paper.

References

- Button KS, Ioannidis JP, Mokrysz C, Nosek BA, Flint J, Robinson ES, Munafò MR** (2013). Power failure: why small sample size undermines the reliability of neuroscience. *Nature Reviews Neuroscience* **14**, 365-376.
- Carp J** (2012). The secret lives of experiments: methods reporting in the fMRI literature. *Neuroimage* **63**, 289-300.
- Ioannidis JP** (2005). Why most published research findings are false. *PLoS Medicine* **2**, e124.
- Kempton MJ, Geddes JR, Ettinger U, Williams SC, Grasby PM** (2008). Meta-analysis, database, and meta-regression of 98 structural imaging studies in bipolar disorder. *Archives of General Psychiatry* **65**, 1017-1032.
- Kempton MJ, Salvador Z, Munafò MR, Geddes JR, Simmons A, Frangou S, Williams SC** (2011). Structural neuroimaging studies in major depressive disorder. Meta-analysis and comparison with bipolar disorder. *Archives of General Psychiatry* **68**, 675-690.
- Park IU, Peacey MW, Munafò MR** (2013). Modelling the effects of subjective and objective decision making in scientific peer review. *Nature* **506**, 93-96.
- Simmons JP, Nelson LD, Simonsohn U** (2011). False-positive psychology: undisclosed flexibility in data collection and analysis allows presenting anything as significant. *Psychological Science* **22**, 1359-1366.
- Simonsohn U** (2013). Just post it: the lesson from two cases of fabricated data detected by statistics alone. *Psychological Science* **24**, 1875-1888.
- Smulders YM** (2013). A two-step manuscript submission process can reduce publication bias. *Journal of Clinical Epidemiology* **66**, 946-947.

Figure 1. Association of Year of Publication with Number of Participants and Number of Measures in Structural Imaging Studies of Major Depression and Bipolar Disorder.

In studies of bipolar disorder (top panel) year of publication is not associated with number of participants ($R^2 = 0.01$, $P = 0.23$) but is associated with number of measures collected ($R^2 = 0.07$, $P = 0.001$). However, in studies of major depression (bottom panel) year of publication is associated with number of participants ($R^2 = 0.04$, $P = 0.001$) but not with number of measures collected ($R^2 = 0.01$, $P = 0.21$).

Analytical Flexibility in Imaging Studies

